



Original article

Effects of sensory motor training of lower limb in sedentary elderly as part of functional autonomy



A.A. Barbosa Rezende^a, E. Fernandes de Miranda^{a,b,*}, H. Souza Ramalho^a, J.D. Borges Da Silva^a, S.D. Silva Carlotto Herrera^a, G. Rossone Reis^a, E.H. Martin Dantas^{b,c}

^a Centro Universitário UnirG, Gurupi-TO, Brazil

^b Laboratory of Biosciences of Human Motricity (LABIMH) of the Federal University of Rio de Janeiro State (UNIRIO), Rio de Janeiro, Brazil

^c University Tiradentes (UNIT), Brazil

ARTICLE INFO

Article history:

Received 7 September 2013

Accepted 28 May 2014

Keywords:

Motor activity

Elderly

Limited mobility

ABSTRACT

Objective: Assess the effects of sensorimotor training on the functional autonomy levels of sedentary elderly people.

Methods: The sample was composed of 30 women (sedentary, functionally independent, $\chi = 65.03 \pm 4.93$ year), divided randomly into two groups: control (CG, $n = 15$) and experimental (EG, $n = 15$). Tests developed by the Latin American Development for the Elderly Group (GDLAM) were used to evaluate autonomy. The EG was submitted to a sensorimotor program three times a week, involving 24 thirty-minute sessions. Data were processed using the SPSS program, with a 5% significance level.

Results: The EG showed a significant reduction in execution time in all the tasks, while the CG only obtained a significantly better score on the 10-M walk test. Although the EG improved in the GDLAM index ($\Delta\% = -30, 3\%$, $p = 0.0001$), it remained at a “weak” level (GDLAM index > 27.42).

Conclusion: The results demonstrate that sensorimotor training reduces the execution time of tasks, as well as improving functional performance in activities of daily living.

© 2013 Consejería de Educación, Cultura y Deporte de la Junta de Andalucía. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Efectos del entrenamiento sensorio motriz del miembro inferior en ancianos sedentarios como parte de su autonomía funcional

RESUMEN

Objetivo: Evaluar los efectos del entrenamiento sensorio motriz en los niveles de autonomía funcional del anciano sedentario.

Método: La muestra se compuso por 30 mujeres (sedentarias, independientes funcionalmente, $65,03 \pm 4,93$ años) divididas al azar en 2 grupos: control (CG, $n = 15$) y experimental (EG, $n = 15$). Para la evaluación de la autonomía se utilizaron las pruebas del Grupo del Desarrollo Latinoamericano para la Madurez (GDLAM) antes y después de la intervención. El EG fue sometido a un programa de entrenamiento sensorio motriz, 3 veces a la semana, la duración de cada sesión fue de 30 minutos y realizaron 24 sesiones. Los datos se procesaron con el programa SPSS, versión 19, adoptando nivel de significación de la $p < 0,05$.

Resultados: En el grupo experimental se constató una reducción significativa del tiempo de ejecución en todas las tareas, mientras el CG solo obtuvo resultado significativamente mejor en la prueba de caminata de 10 m. En el índice general de GDLAM, el grupo experimental presentó mejora ($\Delta\% = -30, 3\%$, $p = 0,0001$), pero se mantuvo en un nivel «débil» (Índice GDLAM $> 27,42$).

Palabras clave:

Actividad motora

Mayores

Limitación de la movilidad

* Corresponding author at: Centro Universitário UnirG, Gurupi-TO, Brazil.

E-mail address: eduardounirg@gmail.com (E. Fernandes de Miranda).

Conclusión: Los resultados muestran que la práctica del entrenamiento sensorio motriz influye de forma positiva en la reducción del tiempo de ejecución de las tareas, lo que se relaciona con el mejor rendimiento funcional de las actividades cotidianas.

© 2013 Consejería de Educación, Cultura y Deporte de la Junta de Andalucía. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Efeitos do treinamento sensório-motor de membros inferiores em idosos sedentários como parte de sua autonomia funcional

R E S U M O

Palavras-chave:
Atividade motora
Idoso
Limitação da mobilidade

Objetivo: Avaliar os efeitos do treinamento sensório-motor sobre os níveis de autonomia funcional de idosos sedentários.

Métodos: A amostra foi composta por 50 mulheres (sedentárias, funcionalmente independentes, $65,03 \pm 4,93$ anos) divididas aleatoriamente em 2 grupos: controle (CG, $n = 15$) e experimental (EG, $n = 15$). Para a avaliação da autonomia funcional foi utilizado o teste Grupo de Desenvolvimento Latino-Americano para a Maturidade (GDLAM) antes e após a intervenção. O grupo EG foi submetido a um programa de treinamento sensório-motor, 3 vezes por semana, com duração de 30 minutos cada sessão, e foram realizadas durante 24 sessões. Os dados foram processados através do programa SPSS, versão 19, adotando nível de significância de $p < 0,05$.

Resultados: No grupo experimental foi constatada uma redução significativa do tempo de execução em todas as tarefas, enquanto que o grupo CG obteve resultados significativamente menores no teste de caminhada de 10 m. No índice geral do GDLAM, o grupo experimental apresentou melhora ($D\% = -30,3\%$, $p = 0,0001$), porém se manteve em nível «fraco» (índice GDLAM $> 27,42$).

Conclusão: Os resultados mostraram que a prática do treinamento sensório-motor influencia de forma positiva na redução do tempo de execução das tarefas, relacionando-se com melhor rendimento funcional das atividades cotidianas.

© 2013 Consejería de Educación, Cultura y Deporte de la Junta de Andalucía. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Mobility is one of the most important aspects of physical function, especially as a prerequisite for executing activities of daily living (ADL) and maintaining independence.¹ Thus, a significant decrease in the gait pattern of elderly individuals hinders the execution of activities such as walking, climbing up and down stairs, moving in and out of bed or a chair, personal hygiene, shopping and remaining active, thereby progressively compromising functional capacity.^{2,3}

The elderly have reduced muscle strength, suffering from negative influence on the ability of sensory discrimination, with consequent loss of motor coordination, balance and proprioception.^{4–6}

Likewise in elderly autonomous, there is a possibility that these defects, over time, may occur, affecting the pattern of movement and decreasing the individual's ability to walk and minimize the risk of falls. Since gait is a complex motor phenomenon, resulting from the interaction of neurological, musculoskeletal, vestibular and somatosensory systems and is one of the most important aspects of physical function and prerequisite for the execution of daily living (ADLs) activities, there may arise disability and functional dependence^{7–9} is important to test the efficacy of interventions aimed to recover and extend the good operational status of the march, aiming to promote functional autonomy of the elderly.

During the aging process these suffer impairment due to various factors, joint, muscle and nerve factors, and also because of balance disorders^{5,10} resulting in limitations in ADLs that negatively affect the self-confidence of the elderly, hampering their execution, due to fear of falling or other physical, psychological or social factors. Moreover, these restrictions may progressively compromise the functional autonomy over time, creating an irreversible cycle, resulting in total dependence or even bed confinement.¹¹

Functional autonomy is defined as follows: autonomy of action, referring to the notion of physical independence; autonomy of will, that is, the possibility of self-determination; and autonomy of thoughts, related to the possibility of judging any situation.^{12,13} Autonomy of action is also referred to as functional autonomy, describing the ability of an elderly individual to independently perform ADLs.¹²

Physical activity has produced positive effects on maintaining functional autonomy and is a preventive factor against functional limitations, that is, the inability to sustain normal daily tasks, in addition to enhancing performance in ADLs.^{14–15}

Several studies have investigated the benefits of physical activity on functional autonomy;^{1,14,15} however, there is a lack in regards to the training of sensorimotor functional activities that simulate displacement influencing ADLs and therefore part of the functional autonomy protocols.

The aim of the present study was to assess the effects of sensorimotor training of the lower limb in sedentary elderly as part of functional autonomy.

Methods

Subjects

Thirty sedentary, autonomous and functionally independent women, aged between 60 and 75 years ($\chi = 65.03 \pm 4.93$) and enrolled in the Physical Activity and Rehabilitation Program for Autonomy in the Elderly (PARPAE) at the Centro Universitário UnirG in Gurupi, Brazil, were randomly selected.

The following exclusion criteria were adopted: the need for walking aids or assistance from another person; suffering from cardiovascular diseases that limit physical exercise; visual acuity

less than 0.3 by the Snellen Table¹⁶; use of orthoses; presence of neurological dysfunctions; orthopedic diseases of the lower limbs that limit ambulation; otoneurological alterations; participation in other physical exercise programs and attending less than 80% of the sessions.

In accordance with Resolution 196/96 and the Helsinki Convention,¹⁷ subjects were informed about study procedures and possible related discomfort, in addition to giving written informed consent.

The project was approved by the Human Research Ethics Committee of Universidade Castelo Branco (UCB/RJ), under protocol no. 0002/2009.

Diagnostic assessment

Study participants were assessed for functional autonomy, according to the Latin American Development for the Elderly Group (GDLAM) protocol,¹² adapted for this study to characterize functional autonomy status in the elderly.

The assessment consisted of five tests, as follows: 10 m Walk (C10m)¹⁸; getting up from a sitting position (LPS)¹⁹; getting up from the prone position (LPDV)²⁰; getting up from a chair and moving around the house (LCLC).²¹ Scores are awarded based on time (seconds) required to execute tasks, such that the lowest time represents the highest level of functional autonomy. These tests are used to calculate the GDLAM index, classifying functional performance as very good < 22.66; 22.66 < good < 24.97; 24.98 < fair < 27.42 and weak > 27.42. The GDLAM index was calculated by normalization between the four autonomy tests. This calculation was elaborated from the following formula:

$$\text{GDLAM index} = \frac{[(\text{C10} + \text{LPS} + \text{LPDV}) \times 2] + \text{LCLC}}{3}$$

For the last test we used a wooden chair with a 50 cm seat-to-floor height and no arm rest; a Cronobio SW 201 water resistant stopwatch (USA); an E.V.A. tatami mat (Brazil); and a Uyustools 50-cm professional measuring tape (USA).

For test the effect of the learning, the day before the intervention was taught until they properly execute the task. In the period preceding it and post intervention, all tests were performed in two trials with an interval of 5 min, recording the shortest time achieved in the execution of each task. These times were compared with autonomy levels of the GDLAM protocol, in order to classify the elderly according to degree of functional autonomy.

Intervention

After the pretest, subjects were randomly divided into the experimental group (EG $n = 15$), submitted to a sensorimotor program and the control group (CG $n = 15$). The EG underwent three 30 min sessions per week for 8 weeks, followed by reassessment.

The CG met weekly for eight weeks and were informed that they would also be submitted to the same sensorial training after the conclusion of the study.

Training was conducted on a previously planned circuit established by PARPAE (Physical Activity and Rehabilitation for Autonomy in the Elderly Program), a community initiative supported by the federal agency FINEP (Research and Project Financing). The circuit was originally created in order to train the gait pattern, so it has tasks that simulate ADLs involving only the lower limbs. It has not been validated, but already there was a publication that demonstrated the effects of this circuit in the gait pattern of sedentary elderly.²² The circuit, denominated Circuit for Assessment and Development of Gait, Balance and Physical Conditioning in the Elderly, was built in a 15.84-meter-long room,

containing eight stations designed for activities using proprioceptive and balance stimuli, as described below and illustrated in Fig. 1.

Station 1 – Walking over 35-cm-wide cylindrical blocks, ranging between 10 and 40 cm in height, using alternate lower limbs and placing both feet on the block before proceeding to the next one. Blocks were arranged in two levels of difficulty: the first level had seven blocks with little variation in height while the second consisted of nine blocks with greater height variation, requiring more exercise intensity.

Station 2 – Stepping over five 0.75-cm-high signal cones, lying flat and placed 1 m apart in a sandbox measuring 11 m × 1.10.

Station 3 – Walking along a curb 10-cm-high, 12-cm-wide and 11-m-long.

Station 4 – Standing on both feet for 20 s with no hand support, in proprioceptive balance.

Station 5 – Moving forwards and sideways to negotiate eight obstacles tied to the ceiling with nylon thread, arranged in two columns measuring 125 cm × 60 cm.

Station 6 – Walking under two 80-cm-long obstacles located 130 cm above the floor, using squatting movements.

Station 7 – Climbing stairs with seven 10-cm-high steps, angled at 45° and equipped with an 85-cm-high handrail, then going down a 3.10-m-long ramp. This procedure was then repeated.

Station 8 – Walking in a straight line over 3 m of mats placed end-to-end on the floor.

Participants were instructed to complete the circuit at a fairly light intensity, corresponding to level 12 on the Borg Scale,²³ which requires approximately 55% of maximum heart rate, ensuring continuity of the activity for the time period established.

They were also advised to use light clothing and no footwear, in order to avoid possible influences by different types of shoes.

On the first day, prior to the experiment, participants were familiarized with the circuit. They were also informed that they could interrupt the maneuver at any time if they felt discomfort. Furthermore, they were instructed not to alter their activities of daily living, in order to reduce possible external influences on the final results.

Statistical analyses

We used descriptive analysis with mean, standard deviation and delta percentage. Sample normality was assessed by the Shapiro-Wilk test. To analyze response variables, the paired Student's *t* or Wilcoxon test was used for intergroup analysis, when appropriate. The value of $p < 0.05$ was adopted for statistical significance, and the SPSS version 19 program was used to assess results.

Results

The scores obtained by the control and experimental groups on the four tests and the GDLAM index, before and after the intervention, are illustrated in Graphs 1, 2 and 3.

The experimental group showed a significant reduction in execution time in all the tasks (Fig. 2), while the control group scored significantly higher only in the 10-M walk (Fig. 3).

With respect to the GDLAM index (Fig. 4), the experimental group decreased from 81.09 to 54.33 ($p = 0.0001$), that it is still considered weak, that is >27.42. The control group increased from 34.75 to 35.49 ($p = 0.3016$).

Discussion

The low functional performance classification observed in the elderly of the present study is in accordance with the premise that




Fig. 1. Illustration of the circuit for assessment and development of gait, balance and physical conditioning in the elderly.

aging, in conjunction with a sedentary lifestyle, provokes a reduction in functional capacity.²⁴ According to a number of studies,^{25,26} the more active elderly preserve functional autonomy for a longer period of time, thereby reducing the risks of functional decline.

Similar data were observed in another study.²⁴ The authors compared the autonomy level between 15 sedentary elderly individuals, that is, who did not engage in any type of regular physical activity and 15 others who walked at a light intensity pace three times a week for 30 min, over a three-month period. The active elderly obtained better results ($p < 0.01$), but when EG values were compared, both groups were classified as weak. In the C10m, LPS, LPDV and LCLC tests, the active elderly exhibited significantly better results ($p < 0.05$) than the other group.^{12,27}

As a strategy to preserve autonomy, physical exercises can be implemented to increase not only endurance and muscle strength, but also balance and mobility, which could reduce risks of falling

and injuries²⁸ and positively impact the performance of activities of daily living.^{29,30}

Physical activity also improves functional capacity, that is, the ability to perform normal activities of daily living with efficiency and independence, and training focusing on functionality is the best means of developing this capacity. According to the principle of specificity, this form of exercise seeks to adapt the human body to normal activities of daily living, using exercises that stimulate proprioception and physical capacities such as strength, flexibility, motor coordination, balance and cardiovascular endurance.^{31,32}

In order to promote successful learning, specific task-directed activities must be intensely performed, evolving to varied exercises and environments. In this training, several distinct strategies must be applied, since most balance demands derive from activities of daily living. For this reason, two aspects must be considered when selecting functional training activities: the patient's safety and level




Fig. 2. Mean scores for the experimental group on the GDLAM protocol. C10m = 10-meter walk; LPS = getting up from a sitting position; LPDV = getting up from the prone position; LCLC = getting up from a chair and moving around the house. *Significantly different in the post-test ($p < 0.01$).




Fig. 3. Mean scores for the control group on the GDLAM protocol. C10m = 10-meter walk; LPS = getting up from a sitting position; LPDV = getting up from the prone position; LCLC = getting up from a chair and moving around the house. *Significantly different in the post-test ($p < 0.05$).




Fig. 4. Mean GDLAM index scores of control and experimental groups. CG – control group, EG – experimental group. *Significantly different in the post-test ($p < 0.01$).

of control, as well as variety in terms of real-life functional tasks and environment.³¹

In the present study, pre- and post-training sensorimotor results in the EG demonstrate task learning, evidenced by the reduced execution time on the tests. Similar studies have shown that a physical activity program involving walking three times a week for 45 min at an intensity between 55% and 75% of maximum heart rate,³³ and another applying the same frequency with a duration of 60 min, over a three-month period, produced shorter execution times on all GDLAM tests, reflecting in improved functional autonomy.³⁴

An experimental study conducted in elderly women (65 ± 6.40 years), separated randomly into an experimental – EG ($n = 31$) and control group – CG ($n = 28$), assessed aerobic endurance, autonomy and quality of life after recreational activities and 1 h walks three times a week for four months. With respect to autonomy, the GDLAM index of the EG exhibited significant improvements when compared to the pre-test and post-test ($p = 0.0001$), in contrast to the CG, where this did not occur ($p = 0.999$).³⁵

Other types of programs also aim at improving functional autonomy. This can be observed in a study of 52 sedentary elderly people, divided into a control (CG) and experimental (EG) group, submitted to a series of 10 Pilates exercises, twice a week for eight weeks. After the training program, it was found that the EG achieved significantly better results on all the tests and in the GDLAM index ($p = 0.035$), and that the CG obtained a significantly higher score ($p = 0.042$) only on the 10 m walk test.³⁶

Yoga has also been shown to improve autonomy when a group that engaged in this activity twice a week for 60 min over a 14-week period, $n = 52$ (66.79 ± 3.30 years) was compared with a control group, $n = 31$ (69.33 ± 4.84 years), resulting in better GDLAM index ($p = 0.0001$).³⁷

Although sensorimotor training provoked significant improvements in task execution time, the GDLAM remained weak. This may be due to the fact that the subjects investigated were elderly and sedentary, in addition to insufficient training duration. Thus, new

studies should be conducted with longer intervention periods and a younger age group than the one used in the present study.

In conclusion, the results show that engaging in sensorimotor training reduces the task execution, which is directly related to improved functional performance in activities of daily living.

Conflicts of interest

The authors declare that they have no conflicts of interest.

References

- Chou CHCL, Hwang CL, Wu YT. Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: a meta-analysis. *Arch Phys Med Rehabil.* 2012;93(2):237–44.
- Faria JC, Machala CC, Dias RC, Dias JMD. The importance of strength training programs for the rehabilitation of muscle function, equilibrium and mobility of the elderly. *Act Fisiat.* 2003;10(3):133–7.
- Ruwer SL, Rossi AC, Simon LF. Balance in the elderly. *Rev Bras Otorrinolaringol.* 2005;71:298–303.
- Verdú E, Ceballos D, Vilches JJ, Navarro X. Influence of aging on peripheral nerve function and regeneration. *J Peripher Nerv Syst.* 2000;5(4):191–208.
- Xu E, Hong Y, Li J, Chan K. Effect of tai chi exercise on proprioception of ankle and knee joints in old people. *Br J Sports Med.* 2004;38(1):50–4.
- Lustri WR, Morelli JGS. Biological aspects of aging. In: Rebelatto JR, Morelli JGS, editors. *Geriatric physical therapy: the practice of care of the elderly.* São Paulo: Manole; 2004.
- Konrad HR, Girardi M, Helfert R. Balance and aging. *Laryngoscope.* 1999;109(9):1454–60.
- Imms FJ, Edholm OG. Studies of gait and mobility in the elderly. *Age Ageing.* 1981;10:147–56.
- Wall JC. Gait. In: Durward BR, Baer GD, Rowe PJ, editors. *Functional human movement: measurement and analysis.* São Paulo: Manole; 2001.
- Matsudo SM, Matsudo VKR, Barros TL. Impact of aging on anthropometric, metabolic and neuromotor variables of physical fitness. *Braz J Mov Sci.* 2000;8(4):21–32.
- Fabrizio SCC, Rodrigues RAP. The elderly perceptions about changes in daily life after fall injuries. *Rev Enferm UERJ.* 2006;14(4):531–7.
- Dantas EHM, Vale RGS. GDLAM'S protocol of functional autonomy evaluation. *Fit Perf J.* 2004;3(3):175–82.
- Benaïm C, Froger J, Compan B, Pelissier J. Évaluation de l'autonomie de la personne âgée. *Ann Readapt Med Phys.* 2005;48(6):336–40.
- Cader S, Pereira ED, Vale RGS, Dantas EHM. Comparación de la fuerza de la musculatura inspiratoria entre mujeres mayores sedentarias y practicas de hidrogimnasia. *Rev Esp Geriatr Gerontol.* 2007;42(5):271–5.
- Cader S, Silva EB, Vale RGS, Bacelar S, Monteiro MD, Dantas EHM. Effect of inspiratory muscles training in maximal inspiratory pressure and functional autonomy of sheltered elderly people. *Motricidade.* 2007;3(1):279–88.
- Falkenstein IA, Cochran DE, Azen SP, Dustin L, Tammewar AM, Kozak I, et al. Comparison of visual acuity in macular degeneration patients measured with Snellen and early treatment diabetic retinopathy study charts. *Ophthalmology.* 2008;115(2):319–23.
- Association WWM. Declaration of Helsinki. Ethical principles for medical research involving human subjects. In: *WMA General Assembly 59th, Seoul, October. 2008.*
- Spilä S, Multanen J, Kallinen M, Era P, Suominen H. Effects of strength and endurance training on isometric muscle strength and walking speed in elderly women. *Acta Physiol Scand.* 1996;156(4):457–64.
- Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol.* 1994;49(2):85–94.
- Alexander NB, Ulbric HJ, Raheja A, Channer D. Rising from the floor in older adults. *J Am Geriatr Soc.* 1997;45(5):564–9.
- Andreotti RA, Okuma SS. Validation of a battery of activities of daily living for elderly physically independent tests. *Rev Paul Educ Fis.* 1999;13(1):46–66.
- Rezende AAB, Silva IL, Beresford H, Batista LA. Evaluation of the effects of a sensory-motor program in gait patterns of elderly. *Fisioter Mov.* 2012;25(2):317–24.
- Borg GAV, Noble BJ. In: Wilmore JH, editor. *Perceived exertion.* Academic Press; 1974.
- Alencar NA, Bezerra JCP, Dantas EHM. Evaluation of the levels of physical activity, functional autonomy and quality of life of elderly women members of the family health program. *Fit Perf J.* 2009;8(5):315–21.
- Bäckmand HM, Kaprio J, Kujala UM, Sarna S. Physical activity, mood and the functioning of daily living a longitudinal study among former elite athletes and referents in middle and old age. *Arch Gerontol Geriatr.* 2009;48(1):1–9.
- Lyra RGS, Ramiro L, Nunes Júnior PC, Santos Filho SD. Comparison of functional autonomy of trained or not-trained elderly. *Rev Bras Fisiol Exerc.* 2010;9(1):16–23.

27. Alencar NA, Souza Júnior JV, Aragão JCB, Ferreira MAF, Dantas EH. Level of physical activity, functional autonomy and quality of life in sedentary and active elderly women. *Fisioter Mov.* 2010;23(3):473–81.
28. Rebelatto JR, Calvo JI, Orejuela JR, Portillo JC. Influence of a long-term physical activity program on hand muscle strength and body flexibility among elderly women. *Rev Bras Fisioter.* 2006;10:127–32.
29. Salminen M, Vahlberg T, Sihvonen S, Sjosten N, Piirtola M, Isoaho R, et al. Effects of risk-based multifactorial fall prevention on postural balance in the community-dwelling aged: a randomized controlled trial. *Arch Geront Geriatr.* 2009;48(1):22–7.
30. Salminen MJ, Vahlberg TJ, Salonoja MT, Aarnio PT, Kivelä SL. Effect of a risk-based multifactorial fall prevention program on the incidence of falls. *J Am Geriatr Soc.* 2009;57(4):612–9.
31. Bonnefoy M, Boutitie F, Mercier C, Gueyffier F, Carre C, Guetemme G, et al. Efficacy of a home-based intervention programme on the physical activity level and functional ability of older people using domestic services: a randomised study. *J Nutr Health Aging.* 2012;16:370–8.
32. Liu-Ambrose T, Nagamatsu LS, Hsu CH, Bolandzadeh N. Emerging concept: 'central benefit model' of exercise in falls prevention. *Br J Sports Med.* 2013;47:115–7.
33. Guimarães AC, Rocha CAQC, Gomes ALM, Cader SA, Dantas EHM. Effects of a program of physical activity on the autonomy level of elderly participants in the program of the family health. *Fit Perf.* 2008;7(1):5–9.
34. Cunha RCL, Fortes MSR, Ferreira MA, Bezerra JCP, Graup S, Nobre GC, et al. Effects of the research of hiking related the profiles of functional autonomy in elderly women supported by the family health program. *Rev Bras Geriatr Gerontol.* 2010;13(2):197–208.
35. Fraga MJ, Cader SA, Ferreira MA, Giani TS, Dantas EHM. Aerobic resistance, functional autonomy and quality of life (QoL) of elderly women impacted by a recreation and walking program. *Arch Gerontol Geriatr.* 2011;52(1):40–3.
36. Rodrigues BGS, Cader AS, Torres NVOB, Oliveira EM, Dantas EHM. Functional autonomy of elderly women practicing Pilates. *Fisioter Pesq.* 2010;17(4):300–5.
37. Gonçalves LC, Vale RGS, Barata NJF, Varejão RV, Dantas EHM. Flexibility, functional autonomy and quality of life (QoL) in elderly yoga practitioners. *Arch Gerontol Geriatr.* 2011;53(2):158–62.